



DOI: 10.18721/JPM.14208

UDC 539.125.4

DIQUARK PARTON DISTRIBUTION FUNCTIONS BASED ON THE LIGHT-FRONT AdS/QCD QUARK-DIQUARK NUCLEON MODEL

B. Rodriguez-Aguilar, Ya.A. Berdnikov

Peter the Great St. Petersburg Polytechnic University,
St. Petersburg, Russian Federation

In the paper, we present a phenomenological unpolarized parton distribution functions (PDFs) for diquarks based on the light front soft-wall AdS/QCD quark-diquark nucleon model. From a probed model consistent with the Drell–Yan–West relation and quark counting rule, we have performed a fit of some free parameters using known phenomenological data of quark PDFs. The model considers the entire set of possible valence diquarks within the nucleon. In our investigation, we focused on the spin-0 ($(ud)_0$), spin-1 ($(ud)_1$) and spin-1 ($(uu)_1$) valence diquarks in the proton. The diquark PDFs obtained can be used in proton-proton collision simulations.

Keywords: diquark, parton distribution function, AdS/QCD, holography

Citation: Rodriguez-Aguilar B., Berdnikov Ya.A., Diquark parton distribution functions based on the light-front AdS/QCD quark-diquark nucleon model, St. Petersburg Polytechnical State University Journal. Physics and Mathematics. 14 (2) (2021) 87–100. DOI: 10.18721/JPM.14208

This is an open access article under the CC BY-NC 4.0 license (<https://creativecommons.org/licenses/by-nc/4.0/>)

ПАРТОННЫЕ ФУНКЦИИ РАСПРЕДЕЛЕНИЯ ДИКВАРКОВ, ОСНОВАННЫЕ НА (АдС/КХД)-МОДЕЛИ НУКЛОНА КВАРК-ДИКВАРК

Б. Родригес-Агила, Я.А. Бердиников

Санкт-Петербургский политехнический университет Петра Великого,
Санкт-Петербург, Российская Федерация

В работе представлены феноменологические неполяризованные партонные функции распределения для дикварков, основанные на модели с мягкой стеной на световом конусе AdS/QCD кварк-дикваркового нуклона. На основе проверенной модели, согласующейся с соотношением Дрелла – Яна – Веста и правилом счета кварков, в работе определен набор параметров с использованием известных феноменологических данных кварковых функций распределения партонов (ФРП). Модель рассматривает все возможные состояния валентных дикварков. В данном исследовании мы остановились на рассмотрении состояний валентных дикварков ud со спинами 0 ($(ud)_0$) и 1 ($(ud)_1$), а также uu со спином 1 ($(uu)_1$) в протоне. Полученные дикварковые ФРП могут быть использованы при моделировании протон-протонных столкновений.

Ключевые слова: дикварк, функция распределения партонов, АдС/КХД, голография

Ссылка при цитировании: Родригес-Агила Б., Бердиников Я.А. Партоны функции распределения дикварков, основанные на (АдС/КХД)-модели нуклона кварк-дикварк // Научно-технические ведомости СПбГПУ. Физико-математические науки. 2021. Т. 14. № 2. С. 87–100. DOI: 10.18721/JPM.14208

Статья открытого доступа, распространяемая по лицензии CC BY-NC 4.0 (<https://creativecommons.org/licenses/by-nc/4.0/>)

Introduction

Since the second half of the last decade of the 20th century, the AdS/CFT correspondence [1] between string theory in anti-de Sitter (AdS) space-time and conformal field theories (CFTs) in physical space-time has been a very active and interesting field of study. Among other things, the wealth of this correspondence stands in the possibility to perform calculations between opposite coupling regimes, strongly coupled theories can be mapped into weakly coupled ones and *vice versa*. CFTs are defined as scale invariant theories, so it is impossible to apply the AdS/CFT correspondence to the quantum chromodynamics (QCD) itself directly.

It is worth noting that this is because the coupling constants change with the renormalization scale μ in QCD that we get the condition under which perturbation theory is valid [2].

Nevertheless, in the strong coupling regime of QCD, the couplings appear to be approximately constant. This is the basis for a light-front holography, an approximation of the AdS/CFT to QCD quantized on the light front (light-front AdS/QCD) [3] that has shown the ability to find analytic solutions in the non-perturbative regime of QCD, like improving predictions of hadron masses and structure properties (see e.g. Ref. [4]).

In this work, we are particularly interested in the fact that light-front AdS/QCD predicts a general form of two particle bound state wave function inside nucleons which cannot be derived simply from valence quarks [4, 5]. This has led to considerable progress in nucleon analytical results considering valence diquarks in their structure, just as light-front wave functions QCD matched with soft-wall AdS/QCD predictions [6 – 8].

Another recent result contemplates the scale evolution of the parton distribution functions (PDFs) for a quark-diquark nucleon model using scale-dependent parameters following the DGLAP (Dokshitzer – Gribov – Lipatov – Altarelli – Parisi) evolution [5], that are consistent with the quark counting rule and Drell – Yan – West relation [9, 10]. Based on these last two results, we have fitted the PDF parameters of the

quark-diquark nucleon model to the available data from NNPDF2.3 QCD + QED NNLO [11] for u and d quarks, in order to get the unpolarized PDFs for the spin-0 $(ud)_0$, spin-1 $(ud)_1$ and spin-1 $(uu)_1$ diquarks. With such parameters available, the diquark PDFs can be used to simulations of proton (and neutron) collisions with participating diquarks.

To consider proton collisions based on a nucleon model with diquark structures inside, it is useful to inspect the properties of the parton model.

The parton model

The cross section for proton-proton collisions can be expressed by the so called improved parton model formula [12]:

$$\sigma_{(P_1, P_2)} = \sum_{i,j} dx_1 dx_2 f_i^1(x_1, \mu) f_j^2(x_2, \mu) \times \hat{\sigma}_{ij}(x_1 P_1, x_2 P_2, \alpha_s(\mu), \mu), \quad (1)$$

where the scripts 1 and 2 are labels to incoming proton beams carried momentum P .

In this scenario, the incoming proton beam is equivalent to a beam made of constituent partons. Typically, these partons are taken as the massless-pointlike elementary particles, quarks and gluons [12], with longitudinal momentum distribution characterized by the parton distribution functions $f_i(x, \mu)$.

This means, given some proton with momentum P , the probability to find in such parton i with momentum between xP and $(x + dx)P$ is precisely $dx f_i(x, \mu)$ being dependent as well of the renormalization scale μ .

While σ represents the parton cross sections, which can be computed with perturbative QCD (pQCD) for sufficiently small running coupling $\alpha_s(\mu)$ [2].

However, due to the fact that partons cannot be observed as free particles, the PDFs cannot be calculated using pQCD. Nowadays, the simplest way to obtain PDFs is fitting observables to experimental data, among other phenomenological tools (see e.g. Refs. [13, 14]).

Nevertheless, in order to work with a parton model using constituent diquarks, we must ex-



expand this picture beyond quarks and gluons. As we mentioned above, recent results from soft-wall AdS/QCD [4, 7] have shown a phenomenological approach to reproduce unpolarized PDFs of quark-diquark nucleons [5].

In the next section we show how this phenomenological approach has been constructed to finally obtain our parameters that allow us to exhibit our diquark PDFs.

The soft-wall light front AdS/QCD quark-diquark nucleon model

In this section we intend to outline how to obtain the PDF of a quark-diquark nucleon model using soft-wall light front holographic QCD (for a more detailed analysis see Ref. [5] and its references, from where this section is heavily based).

To construct such a PDF model, it is assumed that a virtual incoming photon interacts with a massless-valence quark. The other two valence quarks are then forming a spectator diquark. In this way, it is ensured that this model is in accordance with the traditional quark-interacting frameworks, from where it is possible to build reliable properties for the nucleon model, so for diquarks. The diquarks can have then either spin-0 (scalar diquark) or spin-1 (vector diquark).

The nucleon state is represented by a spin-flavor SU(4) symmetry. This implies that the possible states are the isoscalar-scalar diquark singlet state, the isoscalar-vector diquark state and the isovector-vector diquark state. Shortly, the diquark can be either scalar or axial-vector.

For the proton state we can write it as

$$|P; \pm\rangle = C_S |uS^0\rangle^\pm + C_V |uA^0\rangle^\pm + C_{VV} |dA^1\rangle^\pm, \quad (2)$$

where, following the original notation in Ref. [5], S and A represent the scalar and vector diquark having isospin at their superscript; the subscripts in the coefficients denote the isoscalar-scalar (S), the isoscalar-vector state (V) and the isovector-vector state (VV).

For the neutron, the state is given by the isospin symmetry $u \leftrightarrow d$.

Without losing the generality of the model, we

will take the case for the proton, which is what we care about in this work.

Using the light-cone convention $x^\pm = x^0 \pm x^3$ [15] it is convenient to choose a frame where the proton transverse momentum vanishes, denoted as

$$P \equiv \left(P^+, \frac{M^2}{P^+}, \mathbf{0}_\perp \right),$$

where M is the proton mass.

So the momentum of the struck quark can be taken as

$$p \equiv \left(xP^+, \frac{p^2 + |\mathbf{p}_\perp|^2}{xP^+}, \mathbf{p}_\perp \right)$$

and the diquark

$$P_x \equiv ((1-x)P^+, P_x^-, -\mathbf{p}_\perp).$$

We can interpret from this notation that $x = p^+/P^+$ is the longitudinal momentum fraction carried by the struck quark.

Now, we can express the two particle Fock-state expansion.

For total angular momentum projection $J^z = \pm 1/2$ with spin-0 diquark is given by

$$\begin{aligned} |uS\rangle^\pm &= \int \frac{dx d^2 \mathbf{p}_\perp}{2(2\pi)^3 \sqrt{x(1-x)}} \times \\ &\times \left[\psi_+^{\pm(u)}(x, \mathbf{p}_\perp) \left| +\frac{1}{2}s; sP^+, \mathbf{p}_\perp \right\rangle + \right. \\ &\left. + \psi_-^{\pm(u)}(x, \mathbf{p}_\perp) \left| -\frac{1}{2}s; sP^+, \mathbf{p}_\perp \right\rangle \right], \end{aligned} \quad (3)$$

where

$$|\lambda_q \lambda_s; sP^+, \mathbf{p}_\perp\rangle$$

is the two-particle state having struck quark of helicity λ_q and a scalar diquark having helicity $\lambda_s = s$ (spin-0 singlet diquark helicity is denoted by s to distinguish from triplet diquark).

While, the spin-1 diquark state is given by the following expression [16]:

$$\begin{aligned} |\nu A\rangle^\pm = & \int \frac{dx d^2 \mathbf{p}_\perp}{2(2\pi)^3 \sqrt{x(1-x)}} \times \\ & \times \left[\psi_{++}^{\pm(v)}(x, \mathbf{p}_\perp) \left| +\frac{1}{2} + 1; xP^+, \mathbf{p}_\perp \right\rangle + \right. \\ & + \psi_{+-}^{\pm(v)}(x, \mathbf{p}_\perp) \left| -\frac{1}{2} + 1; xP^+, \mathbf{p}_\perp \right\rangle + \\ & + \psi_{+0}^{\pm(v)}(x, \mathbf{p}_\perp) \left| +\frac{1}{2} 0; xP^+, \mathbf{p}_\perp \right\rangle + \\ & + \psi_{-0}^{\pm(v)}(x, \mathbf{p}_\perp) \left| -\frac{1}{2} 0; xP^+, \mathbf{p}_\perp \right\rangle + \\ & + \psi_{+-}^{\pm(v)}(x, \mathbf{p}_\perp) \left| +\frac{1}{2} - 1; xP^+, \mathbf{p}_\perp \right\rangle + \\ & \left. + \psi_{--}^{\pm(v)}(x, \mathbf{p}_\perp) \left| -\frac{1}{2} - 1; xP^+, \mathbf{p}_\perp \right\rangle \right]. \end{aligned} \quad (4)$$

where $|\lambda_q \lambda_D; sP^+, \mathbf{p}_\perp\rangle$ represents a two-particle state with a quark of helicity $\lambda_q = \pm \frac{1}{2}$ and a vector diquark of helicity $\lambda_D = \pm 1, 0$ (triplet). Here $v = u, d$ is a flavor index.

The light-front (LF) wave functions with spin-0 diquark state, $\psi_{\pm}^{\pm(u)}$ at the initial scale μ_0 for $J = \pm \frac{1}{2}$ are given by expressions [8]:

$$J = +\frac{1}{2}: \begin{cases} \psi_+^{+(u)}(x, \mathbf{p}_\perp) = N_s \phi_1^{(u)}(x, \mathbf{p}_\perp), \\ \psi_-^{+(u)}(x, \mathbf{p}_\perp) = \\ = N_s \left(-\frac{p^1 + ip^2}{xM} \right) \phi_2^{(u)}(x, \mathbf{p}_\perp); \end{cases} \quad (5)$$

$$J = -\frac{1}{2}: \begin{cases} \psi_+^{-(u)}(x, \mathbf{p}_\perp) = \\ = N_s \left(\frac{p^1 - ip^2}{xM} \right) \phi_2^{(u)}(x, \mathbf{p}_\perp), \\ \psi_-^{-(u)}(x, \mathbf{p}_\perp) = N_s \phi_1^{(u)}(x, \mathbf{p}_\perp). \end{cases} \quad (6)$$

In a very similar way, for vector diquarks with $J = \pm \frac{1}{2}$ the LF wave functions $\psi_{\pm\pm}^{\pm(v)}$ at the initial scale μ_0 can be written as

$$J = +\frac{1}{2}: \begin{cases} \psi_{++}^{+(v)}(x, \mathbf{p}_\perp) = \\ = N_1^v \sqrt{\frac{2}{3}} \left(\frac{p^1 - ip^2}{xM} \right) \phi_2^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{-+}^{+(v)}(x, \mathbf{p}_\perp) = N_1^v \sqrt{\frac{2}{3}} \phi_1^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{+0}^{+(v)}(x, \mathbf{p}_\perp) = N_0^v \sqrt{\frac{1}{3}} \phi_1^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{-0}^{+(v)}(x, \mathbf{p}_\perp) = \\ = N_0^v \sqrt{\frac{1}{3}} \left(\frac{p^1 + ip^2}{xM} \right) \phi_2^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{+-}^{+(v)}(x, \mathbf{p}_\perp) = 0, \\ \psi_{--}^{+(v)}(x, \mathbf{p}_\perp) = 0; \end{cases} \quad (7)$$

$$J = -\frac{1}{2}: \begin{cases} \psi_{++}^{-(v)}(x, \mathbf{p}_\perp) = 0, \\ \psi_{-+}^{-(v)}(x, \mathbf{p}_\perp) = 0, \\ \psi_{+0}^{-(v)}(x, \mathbf{p}_\perp) = \\ = N_0^v \sqrt{\frac{1}{3}} \left(\frac{p^1 - ip^2}{xM} \right) \phi_2^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{-0}^{-(v)}(x, \mathbf{p}_\perp) = N_0^v \sqrt{\frac{1}{3}} \phi_1^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{+-}^{-(v)}(x, \mathbf{p}_\perp) = N_1^v \sqrt{\frac{2}{3}} \phi_1^{(v)}(x, \mathbf{p}_\perp), \\ \psi_{--}^{-(v)}(x, \mathbf{p}_\perp) = \\ = N_1^v \sqrt{\frac{2}{3}} \left(\frac{p^1 + ip^2}{xM} \right) \phi_2^{(v)}(x, \mathbf{p}_\perp). \end{cases} \quad (8)$$

The LF wave functions $\phi_i^{(v)} (i = 1, 2)$ are the twist-3 LF wave functions. These functions can be derived in light-front QCD and in soft-wall AdS/QCD [4, 17 – 19, 6].

In Ref. [7] a generalized form to $\phi_i^{(v)}$ was proposed by matching the electromagnetic form factors of the nucleon in soft-wall AdS/QCD and light-front QCD, getting that

Table 1
The fitted parameters for nucleon valence u and d quarks at the initial scale μ_0 [5]

Parameter	Value	
	u	d
a_1^v	0.280 ± 0.001	0.5850 ± 0.0003
b_1^v	0.1716 ± 0.0051	0.7000 ± 0.0002
a_2^v	0.84 ± 0.02	$0.9434^{+0.0017}_{-0.0013}$
b_2^v	0.2284 ± 0.0035	$0.6400^{+0.0082}_{-0.0022}$
δ^v	1.0	1.0

Notation: $v = u, d$ – quarks, while $a_i^v, b_i^v, a_2^v, b_2^v, \delta^v$ – parameters defined in Eq. (9).

$$\varphi_i^{(v)}(x, \mathbf{p}_\perp) = \frac{4\pi}{\kappa} \sqrt{\frac{\log(1/x)}{(1-x)}} x^{a_i^v} (1-x)^{b_i^v} \times \\ \times \exp \left[-\delta^v \frac{\mathbf{p}_\perp^2}{2\kappa^2} \frac{\log(1/x)}{(1-x)^2} \right], \quad (9)$$

where κ is a scale parameter coming from the soft-wall AdS/QCD model.

With this information, it is possible to write the Dirac and Pauli form factors for spin-1/2 composite particle systems [20].

In Ref. [21] it was found, by fitting the proton form factors from the soft-wall AdS/QCD model with experimental data [22 – 26], that the best agreement if given with $\kappa = 0.4066$ GeV. Furthermore, in Ref. [5] the flavor form factors for u and d in this light-front diquark model was fitted with experimental data [27, 28], obtaining the value of the parameters $a_i^{(v)}$ and $b_i^{(v)}$ at the initial scale μ_0 (see Table 1).

In the same way, using the Sachs form factors, the coefficients for the quark-diquark nucleon state (2) were obtained in Ref. [5]:

$$C_s^2 = 1.3872, \quad C_\nu^2 = 0.6128, \quad C_{\nu\nu}^2 = 1.0.$$

Besides, the normalized constants N_i were found to be

$$N_S = 2.0191, \quad N_0^{(u)} = 3.2050, \quad N_0^d = 5.9423,$$

$$N_1^{(u)} = 0.9895, \quad N_1^d = 1.1616.$$

Quark-diquark unpolarized PDF evolution

The unpolarized parton distribution function is defined as [8, 5]:

$$f^{(v)}(x, \mu_0) = \frac{1}{2} \int \frac{dz^-}{2(2\pi)} \exp \left(\frac{i p^+ z^-}{2} \right) \times \\ \times \left\langle P; S \left| \bar{\psi}^{(v)}(0) \gamma^+ \psi^{(v)}(z^-) \right| P; S \right\rangle \Big|_{z^+ = z_T = 0}, \quad (10)$$

which depends only on the light-cone momentum fraction $x = p^+/P^+$ where the proton state $|P; S\rangle$ with spin S is given as in Eq. (2).

Indeed, γ^+ is the light-cone representation of the usual γ^μ matrix, detailed definition is found in Ref. [15].

The leading order QCD evolution of the unpolarized PDF is given as the standard DGLAP expansion [29, 30, 5]:

$$\int_0^1 dx x^n f(x, \mu) = \\ = \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{\frac{\gamma_n^{(0)}}{2\beta_0}} \int_0^1 dx x^n f(x, \mu_0), \quad (11)$$

where the anomalous dimension is determined by

$$\gamma_n^{(0)} = -2C_F \left(3 + \frac{2}{(n+1)(n+2)} - 4 \sum_{k=1}^{n+1} \frac{1}{k} \right) \quad (12)$$

and the running coupling constant is given as

$$\alpha_s(\mu) = \frac{4\pi}{\beta_0 \ln\left(\frac{\mu^2}{\Lambda_{\text{QCD}}^2}\right)}. \quad (13)$$

In this work we take $C_F = 4/3$, $\beta_0 = 9$ and $\Lambda_{\text{QCD}} = 0.226 \text{ GeV}$.

The initial scale in most of the works on which ours is based is taken to be $\mu_0 = 0.313 \text{ GeV}$ since it is a value available for pion phenomenology.

Thus, the light-front diquark unpolarized PDFs at scale μ are given by [5]:

$$f^{(S)}(x, \mu) = N_S^2(\mu) \times \\ \times \left[\frac{1}{\delta^u(\mu)} x^{2a_1^u(\mu)} (1-x)^{2b_1^u(\mu)+1} + \right. \\ \left. + x^{2a_2^u(\mu)-2} (1-x)^{2b_2^u(\mu)+3} \times \right. \\ \left. \times \frac{\kappa^2}{(\delta^u(\mu))^2 M^2 \ln(1/x)} \right]; \quad (14)$$

$$f^{(A)}(x, \mu) = \left(\frac{1}{3} N_0^{(v)2}(\mu) + \frac{2}{3} N_1^{(v)2}(\mu) \right) \times \\ \times \left[\frac{1}{\delta^v(\mu)} x^{2a_1^v(\mu)} (1-x)^{2b_1^v(\mu)+1} + \right. \\ \left. + x^{2a_2^v(\mu)-2} (1-x)^{2b_2^v(\mu)+3} \times \right. \\ \left. \times \frac{\kappa^2}{(\delta^v(\mu))^2 M^2 \ln(1/x)} \right]. \quad (15)$$

The parameters a_i^v , b_i^v , δ^v are now dependent on the scale μ such that the relation (11) holds, i. e. [5],

$$a_i^v(\mu) = a_i^v(\mu_0) + A_i^v(\mu), \quad (16)$$

$$b_i^v(\mu) = b_i^v(\mu_0) - \\ - B_i^v(\mu) \frac{4C_F}{\beta_0} \ln\left(\frac{\alpha_s(\mu^2)}{\alpha_s(\mu_0^2)}\right), \quad (17)$$

$$\delta^v(\mu) = \exp\left[\delta_1^v \left(\ln\left(\frac{\mu^2}{\mu_0^2}\right) \right)^{\delta_2^v}\right] \quad (18)$$

where the quantities $A_i^v(\mu)$ and $b_i^v(\mu)$ are defined as

$$\Pi_i^v(\mu) = \alpha_{\Pi,i}^v \mu^{2\beta_{\Pi,i}^v} \left[\ln\left(\frac{\mu^2}{\mu_0^2}\right) \right]^{\gamma_{\Pi,i}^v} \Big|_{i=1,2} \quad (19)$$

for $\Pi = A, B$.

The $a_i^v(\mu)$ and $b_i^v(\mu)$ are the parameters given in Table 1. It should be noted that the parameter δ^v tends to unity while $\mu \rightarrow \mu_0$.

In order to find the evolution parameters $\alpha_{\Pi,i}^v$, $\beta_{\Pi,i}^v$, $\gamma_{\Pi,i}^v$ and δ^v it is useful to write the flavor decomposed PDFs $f^u(x, \mu)$ and $f^d(x, \mu)$. It was well discussed in Ref. [8] that for the relation between quark flavors and diquark states should have a linear behavior with free coefficients to be determinate with experimental data. Indeed, in the same way the proton state (2) has to be consistent with the real world under the same coefficients C_S , C_V and C_{VV} , which was how the flavored form factors were decomposed from the diquarks, and such parameters founded in Ref. [5].

So, the flavor decomposed PDFs are given as

$$f^u(x, \mu) = C_S^2 f^{(S)}(x, \mu) + C_V^2 f^{(V)}(x, \mu), \quad (20)$$

$$f^d(x, \mu) = C_{VV}^2 f^{(VV)}(x, \mu). \quad (21)$$

Then, the flavored PDF $f^v(x, \mu)$ in the light-front quark-diquark model can be written as

$$f^v(x, \mu) = \\ = N^{(v)} \left[\frac{1}{\delta^v(\mu)} x^{2a_1^v(\mu)} (1-x)^{2b_1^v(\mu)+1} + \right. \\ \left. + x^{2a_2^v(\mu)-2} (1-x)^{2b_2^v(\mu)+3} \times \right. \\ \left. \times \frac{\kappa^2}{(\delta^v(\mu))^2 M^2 \ln(1/x)} \right], \quad (22)$$

where

$$N^{(u)} = \left(C_S^2 N_s^2 + C_V^2 \left(\frac{1}{3} N_0^{(u)2} + \frac{2}{3} N_1^{(u)2} \right) \right) \quad (23)$$

and

$$N^{(d)} = \left(C_{VV}^2 \left(\frac{1}{3} N_0^{(d)2} + \frac{2}{3} N_1^{(d)2} \right) \right), \quad (24)$$

for u and d quarks respectively.

In this work, we have followed the fashion of Ref. [5] and we have obtained the values of the evolution parameters by fitting the flavor PDFs (22) with data from NNPDF2.3 QCD + QED NNLO [11].

The fit was performed in gnuplot [31], an open source plotting tool using non-linear least-square theory, taking first a f^v depending on parameters $\Pi_i^v(\mu)$ then getting the evolution parameters $\alpha_{\Pi,i}^v$, $\beta_{\Pi,i}^v$, $\gamma_{\Pi,i}^v$ and δ^v .

The unpolarized PDF data was fitted for 100 equal-spaced data points for different $x \in (0, 1)$ and $\mu^2 = 2, 4, 8, 16, 32, 64, 128, 256 \text{ GeV}^2$.

The fitted parameters for $\alpha_{\Pi,i}^v$, $\beta_{\Pi,i}^v$, $\gamma_{\Pi,i}^v$ are shown in Table 2 while the fitted δ^v being shown

in Table 3.

In appendix A we show the different fits performed for the scales mentioned above.

With this data applied to the PDFs (14) and (15), we have drawn the functions $x \cdot f(x)$ of the isoscalar-scalar diquark and isovector-vector diquark for energy scales $\mu^2 = 10, 10^2, 10^3$ and 10^4 GeV^2 shown in Fig. 1, a, b, c, d respectively. The smooth bands show the case of the scalar diquark, while the checkered bands are for the mentioned vector diquark. It is important to note that

$$\frac{1}{3} N_0^{(u)2} + \frac{2}{3} N_1^{(u)2} \approx N_s^2,$$

from values reported in Ref. [5], so the behavior of the $f^{(S)}$ (isoscalar-scalar) curve and the $f^{(V)}$ (isoscalar-vector) one is very similar.

Table 2
PDF evolution parameters with 95% confidence bounds

$\Pi_i^v(\mu)$	α_i^v	β_i^v	γ_i^v	$\chi^2/d.o.f$
A_1^u	-0.196314 ± 0.002266	-0.197209 ± 0.010210	0.927163 ± 0.036270	0.09
B_1^u	6.48940 ± 0.04592	0.161127 ± 0.006494	-0.910813 ± 0.021850	0.17
A_2^u	-0.441651 ± 0.002674	-0.0389503 ± 0.0058020	0.306214 ± 0.019020	0.995
B_2^u	2.58149 ± 0.26410	-0.0548368 ± 0.0780600	-0.807298 ± 0.277900	1.54
A_1^d	-0.119059 ± 0.002517	-0.124819 ± 0.018800	0.952914 ± 0.060100	0.27
B_1^d	12.84810 ± 0.09134	0.0976609 ± 0.006134	-0.80035 ± 0.01510	0.53
A_2^d	-0.514816 ± 0.000724	-0.001555 ± 0.001244	0.171831 ± 0.003307	0.41
B_2^d	1.10727 ± 0.00703	0.0844447 ± 0.005591	-0.57190 ± 0.01486	0.03

Table 3
PDF evolution parameters δ_1^v and δ_2^v for $v = u, d$

$\delta^v(\mu)$	δ_1^v	δ_2^v	$\chi^2/d.o.f$
δ^u	0.035074 ± 0.03009	0.48314 ± 0.06732	10.50
δ^d	0.406762 ± 0.007024	0.46990 ± 0.01275	3.79

Footnote: in Tabs. 2 and 3 the data was used from NNPDF2.3 QCD+QED NNLO [11].




Fig. 1. Graphs of the $x \cdot f(x)$ functions (diquark PDFs) at different scale energies μ^2 , (GeV) 2 : 10 (a), 10^2 (b), 10^3 (c), 10^4 (d). The cases of the scalar ud_0 (gray bands) and vector uu_1 (black ones) diquarks are shown

Conclusions

The soft-wall light front AdS/QCD has allowed us to construct parton distribution functions (PDFs) for diquarks in agreement with the data obtained for quarks phenomenologically. We have particularly taken data from NNPDF2.3 QCD+QED NNLO [11], but the model can be adapted to desired experimental data with u and d quark PDF information.

Although the uncertainties for the values in $\Pi_i^\gamma(\mu)$ reported here should be still improved, an acceptable fit for the functions (14) and (15) is shown in our parameters in Tabs. 2 and 3 looking at $\chi^2/d.o.f.$.

In general terms, the behavior of diquark PDFs observed in Figs. 1 reveals a similarity to the quark PDFs. Such behavior goes in the sense that as the energy scale increases, a shift to $x = 0$ of the peak of the functions is visible; as well as, while x approaches 1, xf tends to vanish exponentially. This fact can be compared with the re-

sults in Ref. [5], where using the same model with NNPDF21(NNLO) [32] data were fitted the u and d quark PDFs.

The phenomenological diquark PDFs reported here are intended to be tested within the framework of particle collisions.

Especially for us, it is expected to study the effect in the production of hadrons in collision simulations of the AdS/QCD quark-diquark nucleon model taking into account participant diquarks in hard processes.

Acknowledgements

B.R. would like to thank to the Secrearía Nacional de Ciencia y Tecnología (Ref. Grant FINDECY/EDUCA CTi 02-2019) of Guatemala for financial support.

Moreover, many thanks to Anatolii Iu. Egorov (HSEP IPNT Peter the Great SPbPU) for many valuable comments and insightful discussions of the conducted studies.



Appendix

Parameter fitting for PDF evolution from NNPDF2.3 QCD+QED NNLO

The scale evolution of A_i^v and B_i^v is parameterized by α_i^v , β_i^v and γ_i^v . While δ^v is parameterized by δ_1^v and δ_2^v , $f(x, \mu)$ is given by Eq. (22) along with Eqs. (16) – (19). The $f(x, \mu)$ function depending on the parameters A_i^v , B_i^v and δ^v are fitted at 8 different energy scales μ^2 in Table 4 for u quark, while the fitted parameters for d quark are given in Table 5.

Each $\chi^2/d.o.f$ was evaluated for 100 equally-spaced points for different $x \in (0, 1)$. The fitting of the parameters at $\mu^2 = 2, 4, 8, 16, 32, 64, 128$ and 256 GeV^2 are shown in Fig. 2 and 3. The data points are extracted from NNPDF2.3 QCD + QED NNLO [11]. It should be noted that the $\chi^2/d.o.f$ values show that the uncertainty ranges found with the fit are overestimated, this is because for this first instance, uncertainties were not taken from the PDF data of Ref. [11]. An improvement in this fact is expected for future works.




Fig. 2. Plots of δ^u (a) and δ^d (b) parameters vs. energy scales obtained by fitting the data of Table 3 through varying evolution parameters δ_1^v and δ_2^v or u (a) and d (b) quarks

Fitting of the PDF $f_1(x)$ at various energy scales for the u quark

Table 4

μ^2, GeV^2	A_1^u	B_1^u	A_2^u	B_2^u	δ^u	$\chi^2/d.o.f$
2.0	-0.133482 ± 0.027630	9.88657 ± 0.57650	-0.398994 ± 0.008142	2.50897 ± 0.82540	1.16148 ± 0.04635	9.15238e -06
4.0	-0.206116 ± 0.014570	5.97471 ± 0.18860	-0.463197 ± 0.004770	1.64702 ± 0.29730	1.39743 ± 0.03360	2.73019e -06
8.0	-0.257193 ± 0.006954	4.63066 ± 0.06874	-0.508357 ± 0.002519	1.28388 ± 0.10670	1.60899 ± 0.02080	6.44055e -07
16.0	-0.294376 ± 0.002982	3.97296 ± 0.02359	-0.542694 ± 0.001198	1.01137 ± 0.03058	1.80059 ± 0.01118	1.22395e -07
32.0	-0.316551 ± 0.003503	3.63544 ± 0.02254	-0.567223 ± 0.001527	0.774017 ± 0.020070	1.94722 ± 0.01540	1.77367e -07
64.0	-0.325091 ± 0.005228	3.45959 ± 0.02855	0.582866 ± 0.002362	0.620872 ± 0.018460	2.03299 ± 0.02475	4.29130e -07
128.0	-0.325202 ± 0.006335	3.36176 ± 0.03040	-0.592571 ± 0.002883	0.504157 ± 0.019470	2.07403 ± 0.03057	7.16693e -06
256.0	-0.324260 ± 0.006969	3.28934 ± 0.04935	-0.600304 ± 0.003162	0.504157 ± 0.019470	2.10605 ± 0.03384	6.94007e -06




Fig. 3. Plots of A_1^u (a), B_1^u (b), A_2^u (c), B_2^u (d), A_1^d (e), B_1^d (f) and A_2^d (h) parameters vs. energy scales obtained by fitting the data of Tabs. 4, 5 through using Eq. (19) and varying evolution parameters $\alpha_{\Pi,i}^v$, $\beta_{\Pi,i}^v$ and $\gamma_{\Pi,i}^v$ for u (a – d) and d (e – h) quarks

Table 5
Fitting of the PDF $f_i(x)$ at various scales for the d quark

μ^2, GeV^2	A_1^d	B_1^d	A_2^d	B_2^d	δ^d	$\chi^2/d.o.f$
2.0	-0.0726864± ±0.0088830	18.11530± ±0.19740	-0.486242± ±0.001573	1.43562± ±0.08108	1.388050± ±0.008694	1.54472e -06
4.0	-0.142581± ±0.008094	11.07900± ±0.11820	-0.543380± ±0.001465	1.01348± ±0.04888	1.612030± ±0.009794	1.42185e -06
8.0	-0.189572± ±0.007766	8.65931± ±0.09223	-0.582432± ±0.001427	0.860135± ±0.037990	1.79567± ±0.01090	1.35531e -06
16.0	-0.224539± ±0.007578	7.42345± ±0.07912	-0.611889± ±0.001410	0.778533± ±0.032410	1.95505± ±0.01194	1.29922e -06
32.0	-0.251850± ±0.007459	6.67682± ±0.07126	-0.635289± ±0.001402	0.729967± ±0.028960	2.09703± ±0.01291	1.24493e -06
64.0	-0.274124± ±0.007376	6.17642± ±0.06598	-0.654558± ±0.001398	0.697843± ±0.026600	2.22594± ±0.01381	1.19305e -06
128.0	-0.292732± ±0.007316	5.81723± ±0.06219	-0.654558± ±0.001398	0.675414± ±0.024860	2.34434± ±0.01466	1.14381e -06
256.0	-0.308610± ±0.007272	5.54758± ±0.05934	-0.684823± ±0.001397	0.659191± ±0.023530	2.45415± ±0.01547	1.09728e -06

Secrearía Nacional de Ciencia y Tecnología of Guatemala FINDECYT/EDUCA CTi 02-2019.

REFERENCES

1. **Maldacena J.M.**, The large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* 2 (2) (1998) 231–252.
2. **Peskin M.E., Schroeder D.V.**, An introduction to quantum field theory, Addison-Wesley Pub. Co., Reading, USA, 1995.
3. **De Teramond G.F., Brodsky S.J.**, Light-front holography: A first approximation to QCD, *Phys. Rev. Lett.* 102 (8) (2009) 081601.
4. **Brodsky S.J., de Teramond G.F.**, Light-front dynamics and AdS/QCD correspondence: The pion form factor in the space- and time-like regions, *Phys. Rev. D.* 77 (5) (2008) 056007.
5. **Maji T., Chakrabarti D.**, Light front quark-diquark model for the nucleons, *Phys. Rev. D.* 94 (9) (2016) 094020.
6. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.**, Nucleon resonances in AdS/QCD, *Phys. Rev. D.* 87 (1) (2013) 016917.
7. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.**, Light-front quark model consistent with Drell–Yan–West duality and quark counting rules, *Phys. Rev. D.* 89 (5) (2014) 054033 [Erratum: *Phys. Rev. D.* 90 (1) (2014) 019902].
8. **Bacchetta A., Conti F., Radici M.**, Transverse-momentum distributions in a diquark spectator model, *Phys. Rev. D.* 78 (7) (2008) 074010.
9. **Drell S.D., Yan T.M.**, Connection of elastic electromagnetic nucleon form-factors at large Q^2 and deep inelastic structure functions near threshold, *Phys. Rev. Lett.* 24 (4) (1970) 181–185.
10. **West G.B.**, Phenomenological model for the electromagnetic structure of the proton, *Phys. Rev. Lett.* 24 (21) (1970) 1206–1209.
11. **Ball R.D., Bertone V., Carrazza S., et al.**, Parton distributions with QED corrections, *Nucl. Phys. B.* 877 (2) (2013) 290–320.
12. **Nason P.**, Introduction to perturbative QCD, Lecture notes for the 11th Jorge Andre Swieca Summer School on Particle and Fields, January 14–27, 2001, Campos do Jordao, SP, Brazil (2002) 409–486.
13. **Pumplin J., Stump D.R., Huston J., et al.**, New generation of parton distributions with uncertainties from global QCD analysis, *JHEP* 2002 (7) (2002) 12.

14. **Ball R.D., Bertone V., Carrazza S., et al.**, Parton distributions for the LHC run II, *JHEP*. 2015 (4) (2015) 40.
15. **Lepage G.P., Brodsky S.J.**, Exclusive processes in perturbative quantum chromodynamics, *Phys. Rev. D*. 22 (9) (1980) 2157.
16. **Ellis J.R., Hwang D. S., Kotzinian A.**, Sivers asymmetries for inclusive pion and kaon production in deep-inelastic scattering, *Phys. Rev. D*. 80 (7) (2009). 074033.
17. **Brodsky S.J., Cao F.-G., de Teramond G.F.**, Meson transition form factors in light-front holographic QCD, *Phys. Rev. D*. 84 (7) (2011) 075012.
18. **Abidin Z., Carlson C.E.**, Nucleon electromagnetic and gravitational form factors from holography, *Phys. Rev. D*. 79 (11) (2009) 115003.
19. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.**, Chiral symmetry breaking and meson wave functions in soft-wall AdS/QCD, *Phys. Rev. D*. 87 (5) (2013) 056001.
20. **Brodsky S.J., Drell S.D.**, The anomalous magnetic moment and limits on fermion substructure, *Phys. Rev. D*. 22 (8) (1980) 2236.
21. **Chakrabarti D., Mondal C.**, Generalized parton distributions for the proton in AdS/QCD, *Phys. Rev. D*. 88 (7) (2013) 073006.
22. **Gayou O., Wijesooriya K., Afanasev A., et al.**, Measurements of the elastic electromagnetic form-factor ratio $\mu_p G_{Ep}/G_{Mp}$ via polarization transfer, *Phys. Rev. C*. 64 (3) (2001) 038202.
23. **Gayou O., Aniol K.A., Averett T., et al.**, Measurement of G_{Ep}/G_{Mp} in $ep \rightarrow ep$ to $Q^2 = 5.6$ GeV 2 , *Phys. Rev. Lett.* 88 (9) (2002) 092301.
24. **Arrington J., Melnitchouk W., Tjon J.A.**, Global analysis of proton elastic form factor data with two-photon exchange corrections, *Phys. Rev. C*. 76 (3) (2007) 035205.
25. **Punjabi V., Perdrisat C.F., Aniol K.A., et al.**, Proton elastic form-factor ratios to $Q^2 = 3.5$ GeV 2 by polarization transfer, *Phys. Rev. C*. 71 (5) (2005) 055202 [Erratum: *Phys. Rev. C*. 71 (6) (2005) 069902].
26. **Puckett A.J.R., Brash E.J., Jones M.K., et al.**, Recoil polarization measurements of the proton electromagnetic form factor ratio $Q^2 = 8.5$ GeV 2 , *Phys. Rev. Lett.* 104 (24) (2010) 242301.
27. **Cates G.D., de Jager C.W., Riordan S., Wojtsekhowski B.**, Flavor decomposition of the elastic nucleon electromagnetic form factors, *Phys. Rev. Lett.* 106 (25) (2011) 252003.
28. **Diehl M., Kroll P.**, Nucleon form factors, generalized parton distributions and quark angular momentum, *Eur. Phys. J. C*. 73 (4) (2013) 2397.
29. **Altarelli G., Parisi G.**, Asymptotic freedom in parton language, *Nucl. Phys. B*. 126 (2) (1977) 298–318.
30. **Broniowski W., Arriola E.R., Golec-Biernat K.**, Generalized parton distributions of the pion in chiral quark models and their QCD evolution, *Phys. Rev. D*. 77 (3) (2008) 034023.
31. **Williams T., Kelley C.**, gnuplot 5.4. An interactive plotting program. Version 5.4. December, 2020, www.gnuplot.sourceforge.net. Accessed February 28, 2021.
32. **Del Debbio L., Forte S., Latorre J.I., et al.**, Neural network determination of parton distributions: the nonsinglet case, *JHEP*. 2007 (3) (2007) 39.

Received 15.05.2021, accepted 27.05.2021.

THE AUTHORS

RODRIGUEZ-AGUILAR Benjamin

Peter the Great St. Petersburg Polytechnic University

29 Politehnicheskaya St., St. Petersburg, 195251, Russian Federation
rodrigesagilar.l@edu.spbstu.ru

BERDNIKOV Yaroslav A.

Peter the Great St. Petersburg Polytechnic University

29 Politehnicheskaya St., St. Petersburg, 195251, Russian Federation
berdnikov@spbstu.ru



СПИСОК ЛИТЕРАТУРЫ

1. **Maldacena J.M.** The large N limit of superconformal field theories and supergravity // Advances in Theoretical and Mathematical Physics. 1998. Vol. 2. No. 2. Pp. 231–252.
2. **Peskin M.E., Schroeder D.V.** An introduction to quantum field theory. Reading, USA: Addison-Wesley Pub. Co., 1995. 842 p.
3. **De Teramond G.F., Brodsky S.J.** Light-front holography: A first approximation to QCD // Physical Review Letters. 2009. Vol. 102. No. 8. P. 081601.
4. **Brodsky S.J., de Teramond G.F.** Light-front dynamics and AdS/QCD correspondence: The pion form factor in the space- and time-like regions // Physical Review. D. 2008. Vol. 77. No. 5. P. 056007.
5. **Maji T., Chakrabarti D.** Light front quark-diquark model for the nucleons // Physical Review. D. 2016. Vol. 94. No. 9. P. 094020.
6. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.** Nucleon resonances in AdS/QCD // Physical Review. D. 2013. Vol. 87. No. 1. P. 016917.
7. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.** Light-front quark model consistent with Drell – Yan – West duality and quark counting rules // Physical Review. D. 2014. Vol. 89. No. 5. P. 054033 [Erratum: Physical Review. D. 2015. Vol. 92. No. 1. P. 019902].
8. **Bacchetta A., Conti F., Radici M.** Transverse-momentum distributions in a diquark spectator model // Physical Review. D. 2008. Vol. 78. No. 7. P. 074010.
9. **Drell S.D., Yan T.M.** Connection of elastic electromagnetic nucleon form-factors at large Q^2 and deep inelastic structure functions near threshold // Physical Review Letters. 1970. Vol. 24. No. 4. Pp. 181–185.
10. **West G.B.** Phenomenological model for the electromagnetic structure of the proton // Physical Review Letters. 1970. Vol. 24. No. 21. Pp. 1206–1209.
11. **Ball R.D., Bertone V., Carrazza S., Del Debbio L., Forte S., Guffanti A., Hartland N.P., Rojo J.** Parton distributions with QED corrections // Nuclear Physics. B. 2013. Vol. 877. No. 2. Pp. 290–320.
12. **Nason P.** Introduction to perturbative QCD // Lecture notes for the 11th Jorge Andre Swieca Summer School on Particle and Fields. January 14 – 27, 2001, Campos do Jordao, SP, Brazil. 2002. Pp. 409–486.
13. **Pumplin J., Stump D.R., Huston J., Lai H.L., Nadolsky P.M., Tung W.K.** New generation of parton distributions with uncertainties from global QCD analysis // Journal of High Energy Physics. 2002. Vol. 2002. No. 7. P. 12.
14. **Ball R.D., Bertone V., Carrazza S., Deans C.S., Del Debbio L., Forte S., Guffanti A., Hartland N.P., Latorre J.I., Rojo J., Ubiali M.** Parton distributions for the LHC run II // Journal of High Energy Physics. 2015. Vol. 2015. No. 4. P. 40.
15. **Lepage G.P., Brodsky S.J.** Exclusive processes in perturbative quantum chromodynamics // Physical Review. D. 1980. Vol. 22. No. 9. P. 2157.
16. **Ellis J.R., Hwang D.S., Kotzinian A.** Sivers asymmetries for inclusive pion and kaon production in deep-inelastic scattering // Physical Review. D. 2009. Vol. 80. No. 7. P. 074033.
17. **Brodsky S.J., Cao F.-G., de Teramond G.F.** Meson transition form factors in light-front holographic QCD // Physical Review. D. 2011. Vol. 84. No. 7. P. 075012.
18. **Abidin Z., Carlson C.E.** Nucleon electromagnetic and gravitational form factors from holography // Physical Review. D. 2009. Vol. 79. No. 11. P. 115003.
19. **Gutsche T., Lyubovitskij V.E., Schmidt I., Vega A.** Chiral symmetry breaking and meson wave functions in soft-wall AdS/QCD // Physical Review. D. 2013. Vol. 87. No. 5. P. 056001.
20. **Brodsky S.J., Drell S.D.** The anomalous magnetic moment and limits on fermion substructure // Physical Review. D. 1980. Vol. 22. No. 9. P. 2236.
21. **Chakrabarti D., Mondal C.** Generalized parton distributions for the proton in AdS/QCD // Physical Review. D. 2013. Vol. 88. No. 7. P. 073006.
22. **Gayou O., Wijesooriya K., Afanasev A., et al.** Measurements of the elastic electromagnetic form-factor ratio $\mu_p G_{Ep}/G_{Mp}$ via polarization transfer // Physical Review. C. 2001. Vol. 64. No. 3. P. 038202.
23. **Gayou O., Aniol K.A., Averett T., et al.** Meas-

- urement of G_{Mp}/G_{Ep} in $ep \rightarrow ep$ to $Q^2 = 5.6 \text{ GeV}^2//$ Physical Review Letters. 2002. Vol. 88. No. 9. P. 092301.
24. **Arrington J., Melnitchouk W., Tjon J.A.** Global analysis of proton elastic form factor data with two-photon exchange corrections // Physical Review. C. 2007. Vol. 76. No. 3. P. 035205.
25. **Punjabi V., Perdrisat C.F., Aniol K.A., et al.** Proton elastic form-factor ratios to $Q^2 = 3.5 \text{ GeV}^2$ by polarization transfer // Physical Review. C. 2005. Vol. 71. No. 5. P. 055202 [Erratum: Physical Review. C. 2005. Vol. 71. No. 6. P. 069902].
26. **Puckett A.J.R., Brash E.J., Jones M.K., et al.** Recoil polarization measurements of the proton electromagnetic form factor ratio $Q^2 = 8.5 \text{ GeV}^2//$ Physical Review Letters. 2010. Vol. 104. No. 24. P. 242301.
27. **Cates G.D., de Jager C.W., Riordan S., Wojtsekhowski B.** Flavor decomposition of the elastic nucleon electromagnetic form factors // Physical Review Letters. 2011. Vol. 106. No. 25. P. 252003.
28. **Diehl M., Kroll P.** Nucleon form factors, generalized parton distributions and quark angular momentum // European Physical Journal. C. 2013. Vol. 73. No. 4. P. 2397.
29. **Altarelli G., Parisi G.** Asymptotic freedom in parton language // Nuclear Physics. B. 1977. Vol. 126. No. 2. Pp. 298–318.
30. **Broniowski W., Arriola E.R., Golec-Bieniat K.** Generalized parton distributions of the pion in chiral quark models and their QCD evolution // Physical Review. D. 2008. Vol. 77. No. 3. P. 034023.
31. **Williams T., Kelley C.** gnuplot 5.4. An interactive plotting program. Version 5.4. December, 2020 // www.gnuplot.sourceforge.net. Accessed February 28, 2021.
32. **Del Debbio L., Forte S., Latorre J.I., Piccione A., J. Rojo J.** Neural network determination of parton distributions: the nonsinglet case // Journal of High Energy Physics. 2007. Vol. 2007. No. 3. P. 39.

Статья поступила в редакцию 15.05.2021, принята к публикации 27.05.2021.

СВЕДЕНИЯ ОБ АВТОРАХ

РОДРИГЕС-АГИЛАР Бенджамин – студент магистратуры Высшей инженерно-физической школы Санкт-Петербургского политехнического университета Петра Великого, Санкт-Петербург, Российская Федерация.

195251, Российская Федерация, г. Санкт-Петербург, Политехническая ул., 29
rodrigesagilar.l@edu.spbstu.ru

БЕРДНИКОВ Ярослав Александрович – доктор физико-математических наук, профессор Высшей инженерно-физической школы Санкт-Петербургского политехнического университета Петра Великого, Санкт-Петербург, Российская Федерация.

195251, Российская Федерация, г. Санкт-Петербург, Политехническая ул., 29
berdnikov@spbstu.ru